

Neutrino Indirect Detection of Neutralino Dark Matter in the CMSSM.

V. Bertin ¹, E. Nezri ^{1 2}, J. Orloff ²

¹ Centre de Physique des Particules de Marseille
 IN2P3-CNRS, Université de la Méditerranée, F-13288 Marseille Cedex 09

² Laboratoire de Physique Corpusculaire de Clermont-Ferrand
 IN2P3-CNRS, Université Blaise Pascal, F-63177 Aubiere Cedex

email : bertin@cppm.in2p3.fr, nezri@in2p3.fr, orloff@in2p3.fr

Abstract

We study potential signals of neutralino dark matter indirect detection by neutrino telescopes in a wide range of CMSSM parameters. We also compare with direct detection potential signals taking into account in both cases present and future experiment sensitivities. Only models with neutralino annihilation into gauge bosons can satisfy cosmological constraints and current neutrino indirect detection sensitivities. For both direct and indirect detection, only next generation experiments will be able to really test this kind of models.

1 Introduction

Our present understanding of the universe is described in the framework of general relativity and cosmology. The densities of its components are related by [1]:

$$\Omega(a) - 1 = \frac{\Omega_{tot} - 1}{1 - \Omega_{tot} + \Omega_{\Lambda}a^2 + \Omega_{mat}a^{-1} + \Omega_{rel}a^{-2}}, \quad (1)$$

where a is the scale factor. This equation and current experimental results suggest and focus on a flat universe with the following parameters [2]:

Cosmological constant:	$\Omega_{\Lambda} = 0.7 \pm 0.1$
Matter:	$\Omega_{mat} = 0.3 \pm 0.1$
baryonic matter:	$\Omega_b = 0.04 \pm 0.01$; including $\Omega_{vis} \lesssim 0.01$
cold dark matter:	$\Omega_{DM} = 0.26 \pm 0.1$
Relativistic components:	$0.01 \lesssim \Omega_{rel} \lesssim 0.05$
neutrinos:	$0.01 \lesssim \Omega_{\nu} \lesssim 0.05$
photons:	$\Omega_{\gamma} = 4.8_{-0.9}^{+1.3} \times 10^{-5}$
Hubble's constant:	$h \equiv H_0/100 \text{ km}^{-1} \text{ s}^{-1} \text{ Mpc}^{-1} = 0.72 \pm 0.08$

In the framework of the Minimal Supersymmetric Standard Model (MSSM) [3, 4, 5, 6, 7], the lightest supersymmetric particle (LSP) is typically the lightest of the neutralinos $\chi_1 (\equiv \chi)$, χ_2 , χ_3 , χ_4 , the mass eigenstates of the neutral gauge and Higgs boson superpartners ($\tilde{B}$, $\tilde{W}^3$, $\tilde{H}_d^0$, $\tilde{H}_u^0$). In the rest of this paper, the LSP is thus assumed to be the lightest neutralino χ_1 and is called more generically χ . Assuming R-parity conservation ($R \equiv (-1)^{B+L+2S}$), the neutralino is a good stable candidate for cold dark matter [8]. In this context, all sparticles produced after the big-bang give a neutralino in their decay chain, leading to a relic bath of χ in the present universe. These neutralinos could be observed via direct detection (χ interaction with a nucleus of a detector), or indirect detection of their annihilation products (ν , γ , $\bar{p}$, $\bar{D}$, e^+). We will mainly focus in this paper on ν indirect detection. The relic neutralinos are gravitationally captured by massive astrophysical bodies and accumulated at the centre of these objects by successive elastic scatterings on their nuclei. The captured neutralino population annihilates and gives rise to neutrino fluxes which could be detectable in neutrino telescopes like Amanda/Icecube, Antares, Baikal.

2 Neutralino in the CMSSM

In the MSSM, the mass matrix of neutralinos in the basis $(-i\tilde{B}, -i\tilde{W}^3, \tilde{H}_d^0, \tilde{H}_u^0)$ is

$$M_{\chi, \chi_2, \chi_3, \chi_4} = \begin{pmatrix} M_1 & 0 & -m_Z \cos \beta \sin \theta_W & m_Z \sin \beta \sin \theta_W \\ 0 & M_2 & m_Z \cos \beta \cos \theta_W & -m_Z \sin \beta \cos \theta_W \\ -m_Z \cos \beta \sin \theta_W & m_Z \cos \beta \cos \theta_W & 0 & -\mu \\ m_Z \sin \beta \sin \theta_W & -m_Z \sin \beta \cos \theta_W & -\mu & 0 \end{pmatrix} \quad (2)$$

where M_1, M_2 are mass term of $U(1)$ and $SU(2)$ gaugino fields, μ is the higgsino “mass” parameter and $\tan \beta = \langle H_u \rangle / \langle H_d \rangle$ is the ratio of the neutral Higgs vacuum expectation values. The neutralino composition is:

$$\chi = N_1 \tilde{b} + N_2 \tilde{W}^3 + N_3 \tilde{H}_d^0 + N_4 \tilde{H}_u^0 \quad (3)$$

and we define its gaugino fraction as $g_{fac} = |N_1|^2 + |N_2|^2$.

In this model, the introduction of soft terms in the Lagrangian breaks explicitly supersymmetry, leading to a low energy effective theory with 106 parameters. The MSSM is therefore not very predictive, and a non biased exploration of its parameter space is not possible. As a first step, we will therefore as usual concentrate on gravity-mediated supersymmetry breaking in supergravity [9] inspired models, with Grand Unification of soft terms at $E_{GUT} \sim 2.10^{16}$ GeV parameterized by m_0 (common scalar mass), $m_{1/2}$ (common gaugino mass) and A_0 (common trilinear term). Together with $\tan \beta$ and $sgn(\mu)$, these define a 5 parameters constraint MSSM (CMSSM) or mSugra model [10, 11, 12], from which the 106 parameters can be deduced through renormalization group equations (RGE).

Due to the large top Yukawa coupling, renormalization group evolution can drive $m_{H_u}^2|_{Q_{EWSB}}$ and/or $m_{H_d}^2|_{Q_{EWSB}}$ to negative values, so that the electroweak symmetry breaking (EWSB) originates purely in quantum corrections, which realizes radiative electroweak symmetry breaking. Minimization of the scalar potential at the electroweak breaking scale Q_{EWSB} yields the condition:

$$\frac{1}{2}m_Z^2 = \frac{m_{H_d}^2|_{Q_{EWSB}} - m_{H_u}^2|_{Q_{EWSB}} \tan^2 \beta}{\tan^2 \beta - 1} - \mu^2|_{Q_{EWSB}} \underset{\tan \beta \gtrsim 5}{\sim} -m_{H_u}^2|_{Q_{EWSB}} - \mu^2|_{Q_{EWSB}}, \quad (4)$$

where usually Q_{EWSB} is taken as $\sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ [13] to minimize one loop corrections.

Such mSugra models offer the theoretical advantage over generic MSSM models that problems such as Landau poles, charge and color breaking (CCB) minima are partially addressed when dealing with RGE.

In mSugra, the lightest neutralino can exhibit two different natures, depending on the input parameter values [14, 15]:

- an almost pure bino-like neutralino for low m_0 , as the RGE drive $M_1|_{Q_{EWSB}} \simeq 0.41 M_1|_{GUT} = 0.41 m_{1/2} \ll |\mu|_{Q_{EWSB}}$ and $M_2|_{Q_{EWSB}} \simeq 0.83 M_1|_{GUT} = 0.83 m_{1/2} \ll |\mu|_{Q_{EWSB}}$;
- for $m_0 \gtrsim 1000$ GeV, the neutralino picks up some higgsino mixing for $\tan \beta \gtrsim 5$ as the increase of m_0 drives $m_{H_u}^2$ to less negative values so that both $|m_{H_u}^2|$ and $|\mu|$ (via eq. 4) decrease. One can then have $|\mu|_{Q_{EWSB}} \lesssim M_1|_{Q_{EWSB}}$ depending on $m_{1/2}$. When $|\mu|$ is too small, EWSB cannot be achieved.

In addition, the very low m_0 values can lead to models with tachyonic sfermions (low $m_{1/2}$) or a slepton LSP. These behaviors are shown on figure 1 in the $(m_0, m_{1/2})$ plane. The neutralino mass is $\sim 0.4 m_{1/2}$ but is affected by low μ values at high m_0 . Unless explicitly quoted we use $m_t = 174.3$ GeV.

In this paper, the RGE, SUSY spectrum and potential minimization are derived using the SUSPECT program [16, 17] version 2.005 which includes 2 loops RGE [18], the tadpole method to minimize the scalar potential [19] and all radiative corrections in the mass spectrum [20]. The neutralino relic density, including $\chi\chi^+$ and $\chi\chi_2$ coannihilations, and detection signals are calculated by the DarkSUSY (DS) package [21, 22]¹ which correctly implements s -channel poles and thresholds [23]. Our interface (available from nezri@in2p3.fr, submitted for inclusion in DS) shortcuts the DS spectrum recalculation, to comply with the SUSPECT spectra which have been checked [24] against SOFTSUSY [25].

3 Branching ratios in neutralino annihilation

The leading processes in neutralino annihilation at rest are shown in figure 2, where the last two processes are proportional to the higgsino/wino fraction of χ and $\chi_i^{(+)}$. In mSugra models, one can typically distinguish different regions of annihilation branching ratios in the $(m_0, m_{1/2})$ plane, for almost any values of $\tan \beta (\geq 5)$ (see figure 3).

¹These calculations use a neutralino local density $\rho_\chi = 0.3 \text{ GeV/cm}^3$ and a Maxwellian velocity distribution with $v_0 = 220 \text{ km s}^{-1}$

$$A_0=0; \tan\beta=45; \mu>0$$




Figure 1: a) Gaugino fraction, b) value of μ and c) neutralino mass in the $(m_0, m_{1/2})$ plane. Region where χ is not the LSP or EWSB is not achieved are also indicated.




Figure 2: Leading channels in neutralino annihilation at rest, with the parametric dependence of their amplitude, N_i being defined in equation 3 [8].




Figure 3: Typical regions of neutralino annihilation.

Above top thresholds: *i.e.* $m_\chi \gtrsim m_{top} \Leftrightarrow m_{1/2} \gtrsim 400$ GeV or slightly higher in the mixed higgsino region.

Region I: for low m_0 ($< 500 - 1000$ GeV), the neutralino is almost purely bino-like. Despite a tiny higgsino fraction, its annihilation via the Z exchange amplitude, being proportional to the fermion mass, is surprisingly important in the top channel $\chi\chi \xrightarrow{Z} t\bar{t} \propto (m_t m_\chi / m_Z^2)^2 N_{3(4)}^2$. Another potentially important annihilation process is $\chi\chi \xrightarrow{A} b\bar{b}$, whose amplitude $|A_{bb}^A| \propto \frac{\tan\beta m_b}{m_W} \left(\frac{m_\chi}{m_A}\right)^2 N_1 N_{3(4)}$ is strongly enhanced for

high $\tan\beta$, both from the explicit dependence, and because m_A decreases when $\tan\beta$ increases. In the region of interest, their ratio $|A_{t\bar{t}}^Z/A_{b\bar{b}}^A|$ accordingly drops from $\sim 20 - 50$ for $\tan\beta = 10$ down to $\sim 2 - 5$ for $\tan\beta = 45$. Even taking the smaller $t\bar{t}$ phase space (a factor of 0.7) and the other subleading $b\bar{b}$ processes into account, the $b\bar{b}$ domination thus calls for an explanation on figure 4 and even more on figure 13. The only way to account for this domination is to recall the opposite sign of the $\chi\chi \xrightarrow{Z} t\bar{t}$ and $\chi\chi \xrightarrow{\tilde{t}} t\bar{t}$ amplitudes, thus allowing for a cancellation provided their order of magnitude match, which we checked analytically. The opposite happens for the annihilation into $b\bar{b}$: the Z , A and sbottom exchange all add up, and from the behaviour of the non-negligible $A_{b\bar{b}}^A$ amplitude, we expect in the region I an important total annihilation cross section $\sigma_{\chi-\chi}^A$ for high values of $\tan\beta$.

Region II: when m_0 increases ($< 1500 - 2000$ GeV), m_A increases and, for intermediate $\tan\beta$ values, the $\chi\chi \xrightarrow{A} b\bar{b}$ amplitude decreases. Since sfermion masses also increase, the $\chi\chi \xrightarrow{\tilde{t}} t\bar{t}$ channel no longer cancels $\chi\chi \xrightarrow{Z} t\bar{t}$, which becomes the dominant process. However, for high $\tan\beta$, $\chi\chi \xrightarrow{A} b\bar{b}$ is enhanced and stays dominant. Globally, since all scalar masses have been increased by m_0 , and the higgsino fraction is still small, the neutralino annihilation cross section $\sigma_{\chi-\chi}^A$ is smaller than in region I.

Region III: increasing further m_0 ($> 2000 - 2500$ GeV), for any value of $\tan\beta$, one finally approaches the mixed higgsino-bino region. The $\chi\chi \xrightarrow{Z} t\bar{t}$ channel amplitude $A_{t\bar{t}}^Z \propto (\frac{m_t m_\chi}{m_Z^2}) N_{3(4)}^2$ then dominates all other processes, which are suppressed by the increase of scalar masses. So we are left with a $t\bar{t}$ region parallel to the highest higgsino fraction isocurves. The m_0 value separating this region from the previous $\chi\chi \xrightarrow{A} b\bar{b}$ region depends on $\tan\beta$. All in all, the neutralino annihilation cross section $\sigma_{\chi-\chi}^A$ is strongly enhanced by Z exchange near the no-EWSB boundary.




Figure 4: Dominant branching ratios of the neutralino annihilation in the $(m_0, m_{1/2})$ plane

Between top and W/Z thresholds:

Region IV: since $m_\chi < m_{top}$, the neutralino annihilates almost only into $b\bar{b}$. Even if some $\chi\chi \xrightarrow{\tilde{b}} b\bar{b}$ occurs, the $\chi\chi \xrightarrow{A} b\bar{b}$ amplitude is dominant.

Region V: increasing m_0 (and m_A), $\chi\chi \xrightarrow{A} b\bar{b}$ remains dominant and this process is still quite efficient for high values of $\tan\beta$. For intermediate values of $\tan\beta$, $\chi\chi \xrightarrow{A} b\bar{b}$ and $\chi\chi \xrightarrow{Z} b\bar{b}$ both occur but their amplitudes are small and the total annihilation $\sigma_{\chi-\chi}^A$ is not efficient.

Region VI: increasing further m_0 disfavors scalar exchange, but even small, the higgsino fraction allows $\chi\chi \xrightarrow{\tilde{\chi}_i^+} W^+W^-$ and $\chi\chi \xrightarrow{\tilde{\chi}_i} ZZ$ to dominate and enhance the total annihilation cross section $\sigma_{\chi-\chi}^A$. Again, the m_0 values delimiting the boundary with region V depend on $\tan\beta$ (via m_A).

Below W/Z thresholds:

Region VII: the dominant process is $\chi\chi \rightarrow b\bar{b}$ via A and/or Z exchange, depending on m_0 , $\tan\beta$ and the higgsino fraction.

This analysis is illustrated on figures 4 and 13, showing the four most important branching ratios for $\tan\beta = 45$ and 10 (the omitted process $\chi\chi \xrightarrow{A} \tau\tau$ behaves as $\chi\chi \xrightarrow{A} b\bar{b}$ but with a smaller amplitude due to the m_τ/m_b ratio). It further offers a qualitative understanding of the relic density picture (figure 5).




Figure 5: Neutralino relic density in the $(m_0, m_{1/2})$ plane. Grey area indicates the excluded models by current accelerators constraints as discussed in the text.

The annihilation is efficient for low values of m_0 (depending on $\tan\beta$) and for a mixed neutralino along the no-EWSB boundary. This gives the “V” (or “U”) shape of the relic density profile for large (or small) $\tan\beta$. According to the current cosmological parameter values, we take the neutralino as an interesting cold dark matter candidate if $0.025 < \Omega h^2 < 0.3$. Figure 5 also shows the region in the $(m_0, m_{1/2})$ plane excluded by the experimental constraints from the Particle Data Group 2000 [26] implemented in DarkSusy, that we have updated with:

$m_{\tilde{\chi}_1^+} > 104$ GeV; $m_{\tilde{f}} > 100$ GeV for $\tilde{f} = \tilde{t}_1, \tilde{b}_1, \tilde{l}^\pm, \tilde{\nu}$, $m_{\tilde{g}} > 300$ GeV; $m_{\tilde{q}_{1,2}} > 260$ GeV for $\tilde{q} = \tilde{u}, \tilde{d}, \tilde{s}, \tilde{c}$ and $-6 \times 10^{-10} < a_\mu(SUSY) < 58 \times 10^{-10}$ [27, 28].

As noted in previous studies [29, 17], the muon anomalous moment a_μ and $b \rightarrow s\gamma$ branching ratio constraints strongly favour $\mu > 0$, to which we restrict in what follows. In figure 5, the grey tail at large m_0 is excluded by the chargino bound; it directly bites into the region relevant for neutrino indirect detection. Less relevant is the exclusion of small m_0 and $m_{1/2}$ values, which comes both from the Higgs mass limit and the $b \rightarrow s\gamma$ branching ratio, as calculated by default in DS. The range $BR(b \rightarrow s\gamma) = 1 \rightarrow 4 \times 10^{-4}$ chosen by default in DarkSusy may seem too low in view of latest CLEO and Belle measurements [26]. However the leading order calculation [30] implemented underestimates the SM value to 2.4×10^{-4} , while next to leading order corrections give 3.6×10^{-4} [31] so that this range should roughly correspond to $2.2 \rightarrow 5.2 \times 10^{-4}$, excluding a bit more than the range $2 \rightarrow 5 \times 10^{-4}$ chosen for instance in [17]. We have checked that replacing the implemented Higgs limit [32] by an aggressive version of the latest limit [33] ($m_h > 114$ GeV for all $\sin(\beta - \alpha)$) only excludes a few more points around $(m_0 = 1000, m_{1/2} = 150)$ where neutrino fluxes are beyond reach.

In mSugra, $\chi\chi^+$ and $\chi\chi_2$ coannihilations (included in DS) happen only in the mixed neutralino region, decreasing further the relic density. $\chi\tilde{\tau}$ coannihilation happens for low values of $\tan\beta$, for which there is no mixed region. $\chi\tilde{t}$ coannihilation [34, 17, 35] happens for high values of A_0 . In both cases, sfermion coannihilations (missing in DS) are relevant to lower the relic density in regions of large m_χ , which as we will see, are beyond reach of indirect detection. We therefore do not expect their proper inclusion to change our conclusions.

4 Neutralino-proton cross section: capture rate and direct detection signals

Capture:

If present in the halo, relic neutralinos must accumulate in astrophysical bodies (of mass M_b) like the Sun or the Earth. The capture rate C depends on the neutralino-quark elastic cross section: $\sigma_{\chi-q}$. Neutralinos being Majorana particles, their vectorial interaction vanishes and the allowed interactions are scalar (via $\chi q \xrightarrow{H,h} \chi q$ in t channel and $\chi q \xrightarrow{\bar{q}} \chi q$ in s channel) and axial (via $\chi q \xrightarrow{Z} \chi q$ in t channel and $\chi q \xrightarrow{\bar{q}} \chi q$ in s channel). Depending on the spin content of the nuclei N present in the body, scalar and/or axial interactions are involved. Roughly, $C \sim \frac{\rho_\chi}{v_\chi} \sum_N M_b f_N \frac{\sigma_N}{m_\chi m_N} < v_{esc}^2 >_N F(v_\chi, v_{esc}, m_\chi, m_N)$, where ρ_χ, v_χ are the local neutralino density and velocity, f_N is the density of nucleus N in the body, σ_N the nucleus-neutralino elastic cross section, v_{esc} the escape velocity and F a suppression factor depending on masses and velocity mismatching. The neutralino capture is maximized for $m_\chi \sim m_N$ as $\frac{\sigma_N}{m_\chi m_N} \sim \frac{m_p^2}{m_\chi m_N} \doteq \frac{m_\chi m_N}{(m_\chi + m_N)^2}$, and is much more efficient in the Sun than in the Earth as $M_\odot \gg M_\oplus$.

For the Earth, scalar interactions dominate. By increasing m_0 from its low value region, sfermions and H exchanges first decrease, and the cross-section rises again when approaching the mixed higgsino region (see figure 6a). The capture rate is resonant for $m_\chi \sim 56$ GeV around the iron mass.

For the Sun, the spin of hydrogen allows for axial interaction, which are stronger due to the Z coupling. The latter depends strongly on the neutralino higgsino fraction and is independent of $\tan \beta$, so the cross-section follows the higgsino fraction isocurves as can be seen by comparing figure 6b and 1.




Figure 6: Scalar (a) and axial (b) cross sections for neutralino scattering on proton in pb.

To summarize, due to the large solar mass and the spin dependent neutralino-quark cross section, the storage of neutralinos is much more efficient in the Sun than in the Earth.

Direct detection:

The elastic cross section of a neutralino on a nucleus also depends on $\sigma_{\chi-q}$, the nucleus mass number A and its spin content. Depending on the chosen nuclei target, current and future direct detection experiments are sensitive to the scalar coupling (CDMS (Ge) [36], Edelweiss (Ge) [37, 38], Zeplin (Xe) [39]) or to the axial coupling (MACHe3 (^3He) [40]). Comparison between experiment sensitivities and a set of mSugra models are shown on figure 9a.

5 Neutrino indirect detection

As $\chi\chi \rightarrow \nu\bar{\nu}$ is strongly suppressed by the tiny neutrino mass, neutrino fluxes come from decays of primary annihilation products, with a mean energy $E_\nu \sim \frac{m_\chi}{2}$ to $\frac{m_\chi}{3}$. The most energetic spectra, called “hard” come from neutralino annihilations into WW , ZZ and the less energetic, “soft”, ones comes from $b\bar{b}$ [21]. Muon neutrinos give rise to muon fluxes by charged-current interactions in the Earth. As both the ν charged-current cross section on nuclei and the produced muon range are proportional to E_ν , high energy neutrinos are easier to detect. Considering that the population of captured neutralinos has a velocity below the escape velocity, and therefore neglecting evaporation, the number N_χ of neutralinos in the centre of a massive astrophysical object depends on the balance between capture and annihilation rates: $\dot{N}_\chi = C - C_A N_\chi^2$,

where C_A is the total annihilation cross section $\sigma_{\chi-\chi}^A$ times relative velocity per volume. The annihilation rate at a given time t is then:

$$\Gamma_A = \frac{1}{2} C_A N_\chi^2 = \frac{C}{2} \tanh^2 \sqrt{C C_A t} \quad (5)$$

with $\Gamma_A \approx \frac{C}{2} = cste$ when the neutralino population has reached equilibrium, and $\Gamma_A \approx \frac{1}{2} C^2 C_A t^2$ in the initial collection period (relevant in the Earth). So, when accretion is efficient, the annihilation rate does not depend on annihilation processes, but follows the capture rate C and thus the neutralino-quark elastic cross section.




Figure 7: ν fluxes from the Sun (a), the corresponding μ fluxes with $E_\mu > 5$ GeV threshold (b) and their ratio (c).

For the Sun, neutralinos do not reach complete equilibrium in the whole $(m_0, m_{1/2})$ plane studied, which contrasts with [29] based on Neutdriver [8] and different RGE, probably yielding lighter (pseudo-)scalars and lower μ . We find that the annihilation is on average less efficient (as attested by our smaller region with an acceptable relic density), which results in a filling fraction $\sqrt{C C_A t_\odot}$ smaller by a factor of 10 to 100. This has little effect on $\nu(\mu)$ fluxes for high m_0 or small $m_{1/2} < 600$ GeV values, as equilibrium is nevertheless reached and fluxes follow essentially the higgsino fraction and the spin dependent $\sigma_{\chi-p}^{spin}$ isocurves of figures 1a and 6. For low m_0 however, the equilibrium fluxes would drop with m_0 and C , but the smaller capture rate hinders equilibrium (e.g. $\sqrt{C C_A t_\odot} \sim 1$ for $m_0 = 500$, $m_{1/2} = 800$) and Γ_A feels the increasing annihilation cross-section. This effect is stronger for high $m_{1/2}$ values where the neutralinos are heavier and more bino-like, and where capture is smaller. The effect is also larger when $\tan \beta$ (and thus $\sigma_{\chi-\chi}^A$) is low, as incomplete equilibrium makes $\nu(\mu)$ fluxes sensitive to $\sigma_{\chi-\chi}^A$, making the W and top thresholds more conspicuous.




Figure 8: ν fluxes from the Earth (a), the corresponding μ fluxes $E_\mu > 5$ GeV threshold (b) and their ratio (c).

For the Earth, neutralinos are not in equilibrium. Neutrino fluxes depend both on C^2 and annihilation, giving an enhancement in the low and high m_0 regions where fluxes are boosted by annihilation (see figure 6 and 8). Since $M_\oplus < M_\odot$ and $\sigma_{\chi-p}^{scal} < \sigma_{\chi-p}^{spin}$, the capture rate and ν fluxes from the Earth are much smaller than from the Sun.

Comparing ν fluxes and μ fluxes, the $\nu \rightarrow \mu$ conversion factor increases with $m_{1/2}$ due to the increase of m_χ leading to more energetic neutrinos. This ratio also follows the annihilation final state regions described in section 3 above. Indeed, spectra from WW , ZZ and to a lesser extent $t\bar{t}$ are more energetic than $b\bar{b}$ spectra, so neutrino conversion into muon is more efficient in the mixed bino-higgsino region above W and top thresholds (see figure 7 and 8).




Figure 9: a) Direct detection experiment sensitivities and a set of models in the $(\sigma_{\chi-p}^{scal}, m_\chi)$ plane. The models excluded by current accelerators constraints are not displayed. b) Indirect detection experiment sensitivities in the $(\mu flux_\odot, m_\chi)$ plane with $E_\mu = 5$ GeV threshold of the same sample as a). Dotted, dash-dotted and dashed curves are respectively the Macro, Baksan and Super-Kamiokande upper limits. Solid lines indicate the future Antares and Icecube sensitivities.

Comparison with indirect detection experiment sensitivities: Muon fluxes coming from neutralino annihilation in the Sun are shown in figure 9b. MSugra models predicting a good relic density of neutralinos give neutrino/muon fluxes which can be as high as the current experimental limits (Baksan [41], Macro [42], Super Kamiokande [43]). The 0.1 km² Antares detector will explore further the interesting parameter space [44]. Next generation neutrinos telescopes (Icecube, Antares km³) will be much more efficient to test such models, eg Icecube expected sensitivity $\sim 10^2 \mu \text{ km}^2 \text{ yr}^{-1}$ from the Sun [45].

Indirect vs Direct detection: a high neutralino-proton cross section is efficient in both direct and indirect detection (via capture). The large m_0 mixed higgsino-bino region pointed out in [15, 29] favours both direct and indirect detection due to the enhancement of $\sigma_{\chi-p}$ and $\sigma_{\chi-\chi}^A$. Both enter in the indirect detection which is moreover favoured by the production of more energetic neutrinos in WW , ZZ and $t\bar{t}$ decays, leading to better conversion into muons (figure 10). This mixed region, which has a good relic density, is very attractive for neutrino indirect detection signal.

However, no current experiment is able to really test such models. Figure 11 shows the region of the $(m_0, m_{1/2})$ plane which can be explored by future direct detection experiments and neutrino telescopes.

6 Variations

In this section, we discuss the robustness of the indirect detection picture described above, with respect to variations in Suspect 2.002 of 1) mSugra input parameters like A_0 or $\tan\beta$, 2) experimental uncertainties on the top quark mass and 3) theoretical uncertainties on the scale of electroweak symmetry breaking Q_{EWSB}

$$A_0=0 ; \tan\beta=45 ; \mu > 0$$




Figure 10: Comparison between μ fluxes from the Sun and neutralino-proton cross sections: a) scalar, b) spin dependent

A₀: Varying A_0 away from 0 does not change too much the analysis above. Annihilation branching ratio regions, as well as the no-EWSB boundary and slope can be slightly displaced. For moderate $\tan\beta \sim 20$, quite large and negative values of A_0 (eg -800 GeV) can enhance $\tau^+\tau^-$ annihilation channel along the $\tilde{\tau}$ LSP region (*i.e.* low m_0) due to splitting in the $\tilde{\tau}$ mass matrix. For such values, the $\tilde{\tau}\chi$ coannihilation [46] could also affect the relic density, but in this region, neutrino fluxes are too small to be detected anyway (see figure 12). The same happens with the $t\bar{t}$ channel for $400 < m_{1/2} < 600$ and $m_0 < 700$ ($\sim |A_0|$). The $\tilde{t}\chi$ [34, 17, 35] coannihilation can also happen for $A_0 \sim -2000$ GeV, but the usual cosmologically interesting region is reduced (due to the negative squared mass arising in sfermion matrices at low m_0 and because the mixed region is pushed to higher values of m_0). In addition, this coannihilation region produces neutrino fluxes from the Sun around $10^{-1} \mu \text{ km}^{-2} \text{ yr}^{-1}$, too low to be detected. In view of the latest results on cosmological parameters, figure 12a shows models with $0.03 < \Omega h^2 < 0.3$ in the $(m_0, m_{1/2})$ plane for different values of A_0 , as well as models giving μ fluxes from the Sun larger than 10^2 (b) and $10^3 \mu \text{ km}^{-2} \text{ yr}^{-1}$ (c). Latter models are confined in the mixed higgsino-bino region. In addition, only those models with $m_W < m_\chi < m_t$ giving a hard spectrum of neutrinos via $\chi\chi \xrightarrow{\tilde{\chi}_i^+} W^+W^-$ and $\chi\chi \xrightarrow{\tilde{\chi}_i} ZZ$ can simultaneously satisfy $0.1 < \Omega h^2 < 0.3$, while yielding a high muon flux.

$\tan\beta$: the branching ratio picture does not qualitatively change for different values of $\tan\beta$. Low values of $\tan\beta$ can add a $\chi\chi \rightarrow t\bar{t}$ region for low m_0 and open a $\tilde{\tau}\chi$ coannihilation [46] region along the $\tilde{\tau}$ LSP region, but μ fluxes are small and such values of $\tan\beta$ reduce the mixed neutralino region which is the most interesting for indirect detection. As discussed previously, the boundaries of branching ratio regions move with $\tan\beta$ (compare figure 13 and 4). When $\tan\beta$ grows, the cosmologically motivated region is wider due to the annihilation enhancement via A exchange (in regions I/II and IV/V). Applying the conservative cut $0.03 < \Omega h^2 < 0.3$, all values of $\tan\beta$ provide interesting models for neutrino indirect detection (see figure 14). For too large values of $\tan\beta$, the $b\bar{b}$ region is extended and leads to soft neutrino spectra. In this case it becomes more difficult to conciliate a tighter limit on the relic density with current experimental sensitivities. Values of $\tan\beta \sim 10 \rightarrow 35$ give more models satisfying both relic density $0.1 < \Omega h^2 < 0.3$ and high ν/μ fluxes.

EWSB scale: Q_{EWSB} is the scale at which the running of the soft SUSY breaking terms is frozen and the minimization of the scalar potential is achieved. When the full potential is minimized within a multi-scale RG treatment, Q_{EWSB} should not affect physics. However, the one-loop single scale effective potential does depend on Q_{EWSB} , and the tree-level relations (equation 4) only hold in terms of running parameters at Q_{EWSB} ; if Q_{EWSB} is chosen $\sim \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ [13], the potential can be minimized perturbatively. Varying Q_{EWSB} away from this value is therefore not physical, but gives a hint on the theoretical uncertainties associated with a single scale RG and potential minimization approach. When Q_{EWSB} is lowered, the RG running drives μ to smaller values. This leads to a wider region of the $(m_0, m_{1/2})$ plane where EWSB cannot be achieved. In the remaining allowed region, the lower values of μ slightly increases the higgsino




Figure 11: a) ν telescopes sensitivities on μ fluxes from the Sun and b) direct detection experiments sensitivities in the $(m_0, m_{1/2})$ plane.

fraction of the neutralino and the annihilation cross-section, giving more cosmologically acceptable models. Figure 15 and 16 show the relic density and μ fluxes from the Sun and the Earth in the $(m_0, m_{1/2})$ plane for $Q_{EWSB} = \frac{1}{2}\sqrt{m_{\tilde{t}_1}m_{\tilde{t}_1}}$ and $\frac{1}{5}\sqrt{m_{\tilde{t}_1}m_{\tilde{t}_2}}$. For $Q_{EWSB} = \frac{1}{5}\sqrt{m_{\tilde{t}_1}m_{\tilde{t}_2}}$, one sees around $m_0 = m_{1/2}$ the $\chi\chi \xrightarrow{A} b\bar{b} s$ -channel annihilation pole which would otherwise show up for higher values of $\tan\beta \gtrsim 55$ [17]. It affects significantly the fluxes from the Earth, where capture and annihilation are not in equilibrium figure 16c. The higher higgsino fraction also enhances the elastic cross section $\sigma_{\chi-q}$, but the stronger annihilation reduces the number of models which are both interesting from the cosmological point of view and in terms of Sun muon fluxes (figure 19c). As before, muon fluxes from the Earth are much too weak to be detected. Increasing Q_{EWSB} leads to the opposite effect: higher values of $|\mu|$ and reduced cosmological favoured region.

Top mass: the current experimental value is $m_t = 174.3 \pm 5.1$ GeV [26]. Our default choice up to now has been to adopt the central value. Lowering m_t implies that the top Yukawa coupling Y_t is smaller so that the RG decreasing of $m_{H_u}^2$ is less efficient and the region where EWSB does not occur is wider (see eq. 4, figure 17 and 18). Nevertheless the mixed higgsino region is larger favouring annihilation and capture, but there are no additional models combining a large ν/μ flux with an acceptable relic density (figure 17, 18 and 19). Increasing the top mass leads to the opposite effect, which is less interesting for our study. It should also be noticed that varying Q_{EWSB} and/or m_t allows for heavier neutralino within the fixed Ωh^2 interval.

7 Conclusion

In this paper, we have analysed, within the framework of the constrained Minimal Supersymmetric Standard Model (*a.k.a.* mSugra), the possible indirect detection of neutralino dark matter through the neutrinos produced by its annihilation in the Earth and the Sun. In particular, we have presented an original study of the relative weight of various annihilation channels crucial to indirect detection. Fixing the halo dark matter density to $\rho_\chi = 0.3$ GeV/cm³ and allowing for the widest range of cosmological relic density $0.03 < \Omega h^2 < 0.3$, we find that a neutrino signal from the centre of the Earth is beyond reach of the present and even future neutrino telescopes.

For the larger Sun, we find that a neutrino signal may be found, but only in the large m_0 "focus point" region [15, 29] where the neutralino has a larger higgsino component which increases both neutralino-nucleon elastic scattering and neutralino annihilation cross-sections. Since in this region, both are related by s - and t -channel crossing (exactly above $t\bar{t}$ threshold and approximately below), any parameter change reducing the relic density inevitably increases the neutrino flux. When conversion into muons is taken into account with a 5 GeV threshold on the muon energy, the only models providing both a relevant (but rather low) relic density, and fluxes around the *current* indirect detection sensitivity ($10^3 \mu \text{ km}^{-2} \text{ yr}^{-1}$) are those with $m_\chi < m_t$ and

$\tan(\beta)=35 ; \mu > 0$




Figure 12: a) Cosmologically favoured models for different values of A_0 in the $(m_0, m_{1/2})$ plane; subset of these models generating a μ flux from the Sun larger than b) $10^2 \text{ km}^{-2} \text{ yr}^{-1}$; c) $10^3 \text{ km}^{-2} \text{ yr}^{-1}$.

$A_0=0 ; \tan(\beta)=10 ; \mu > 0$




Figure 13: Branching ratio of the neutralino annihilation in the $(m_0, m_{1/2})$ plane (same as figure 4 for $\tan \beta = 10$).

$$A_0=0 ; \mu > 0$$




Figure 14: a) Cosmologically favoured models for different values of $\tan\beta$ in the $(m_0, m_{1/2})$ plane; subset of these models generating a μ flux from the Sun larger than b) $10^2 \text{ km}^{-2} \text{ yr}^{-1}$; c) $10^3 \text{ km}^{-2} \text{ yr}^{-1}$.

$$A_0=0 ; \tan(\beta)=45 ; \mu > 0 ; m_{\text{top}}=174.3 \text{ GeV} ; Q_{\text{EWSB}}=1/2 \sqrt{m_{\text{stop1}} m_{\text{stop2}}}$$




Figure 15: a) Relic density, μ fluxes from the Sun (b) and the Earth (c) in the $(m_0, m_{1/2})$ plane for $Q_{\text{EWSB}} = \frac{1}{2} \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ and $m_t = 174.3 \text{ GeV}$.

$\chi\chi \xrightarrow{\chi_i^+} W^+W^-$ or $\chi\chi \xrightarrow{\chi_i} ZZ$ as dominant annihilation processes. Relaxing the tight mSugra constraints in specific directions could favorise such channels [47]. The next generation of neutrino telescopes (with a km^3 size) will be much more efficient to probe mSugra models, especially for $m_\chi > m_t$. If a neutrino signal is found in this framework, then neutralino dark matter should also be accessible to future direct detection experiments, and a chargino with $m_{\chi^\pm} < 350 \text{ GeV}$ should be found in accelerators.

To conclude, a comparison with previous related work is in order. Earlier studies [8, 48] of neutralino indirect detection were performed in the unconstrained MSSM, where μ is a free parameter, and mixed neutralinos occur rather easily. Taking an mSugra slice in this huge parameter space obviously introduces many correlations, for instance between the neutrino flux and the relic density or $\sigma_{\chi-p}^{\text{spin}}$ (and even $\sigma_{\chi-p}^{\text{scal}}$). The effect of this slicing in the $(m_\chi, \sigma_{\chi-p}^{\text{scal}})$ plane was shown in [49] for small $\tan\beta$ and in [50] for large $\tan\beta$. Our figure 9 agrees with this last result, with the addition of the focus point region forming the upper-right cloud. We should note however that a comparison in the $(m_0, m_{1/2})$ plane is more difficult, specially at large $\tan\beta$, because of the theoretical uncertainties in the RGE used to translate these parameters into physical quantities. As an example, for a given value of $\tan\beta$, the slope of the EWSB boundary decreases between

$$A_0=0; \tan(\beta)=45; \mu > 0; m_{\text{top}}=174.3 \text{ GeV}; Q_{\text{EWSB}}=1/5 \sqrt{m_{\text{stop1}} m_{\text{stop2}}}$$




Figure 16: a) Relic density, μ fluxes from the Sun (b) and the Earth (c) in the $(m_0, m_{1/2})$ plane for $Q_{\text{EWSB}} = \frac{1}{5} \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ and $m_t = 174.3 \text{ GeV}$.

$$A_0=0; \tan(\beta)=45; \mu > 0; m_{\text{top}}=170 \text{ GeV}; Q_{\text{EWSB}}=\sqrt{m_{\text{stop1}} m_{\text{stop2}}}$$




Figure 17: a) Relic density, μ fluxes from the Sun (b) and the Earth (c) in the $(m_0, m_{1/2})$ plane for $Q_{\text{EWSB}} = \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ and $m_t = 170 \text{ GeV}$.

$$A_0=0; \tan(\beta)=45; \mu > 0; m_{\text{top}}=170 \text{ GeV}; Q_{\text{EWSB}}=1/2 \sqrt{m_{\text{stop1}} m_{\text{stop2}}}$$




Figure 18: a) Relic density, μ fluxes from the Sun (b) and the Earth (c) in the $(m_0, m_{1/2})$ plane for $Q_{\text{EWSB}} = \frac{1}{2} \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$ for $m_t = 170 \text{ GeV}$.

[29], this work and [50]. As another example, the strong annihilation via s -channel A, H_0 scalars appears above $\tan \beta > 50$ in [51] and [52], $\tan \beta > 35$ in [50], and $\tan \beta > 60$ (or $\tan \beta = 45$, $Q_{\text{EWSB}} = \sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}/5$,

$$A_0=0 ; \tan(\beta)=45 ; \mu > 0$$




Figure 19: a) Dark matter favoured models for different values of Q_{EWSB} and top mass in the $(m_0, m_{1/2})$ plane (a). Colors and symbols correspond to different values of m_t and Q_{EWSB} : e.g. (170 ; 1/2) means $m_t = 170$ GeV and $Q_{EWSB} = \frac{1}{2}\sqrt{m_{\tilde{t}_1} m_{\tilde{t}_2}}$. The models that moreover yield a muon flux from the Sun larger than $> 10^2 \text{ km}^{-2} \text{ yr}^{-1}$ and $> 10^3 \text{ km}^{-2} \text{ yr}^{-1}$ are shown in b) and c).

see figure 16) in the present work using SUSPECT. Finally, we find a muon event rate compatible with [29], and larger than [51]. Part of this difference comes from a higher threshold (25 GeV instead of our 5 GeV). The rest might be attributed to a high sensitivity in the renormalization group equations at large m_0 and $\tan\beta$. It seems [24] that using SOFTSUSY [25] or SUSPECT [16, 17] may be safer than ISASUGRA[53] in this region.

Acknowledgement This work would not have started without the french "GDR supersymetrie". We gratefully thank Jean-Loic Kneur, Charling Tao, and the Antares Neutralino WG for help and stimulating discussions. We must also underline the acute reading and very constructive role of the referee in the final version of this article.

References

- [1] J. A. Peacock. *Cosmological Physics*, chapter 3. Cambridge University Press, 1999.
- [2] C. H. Lineweaver. Cosmological parameters. 2001. talk presented at COSMO-01, Rovaniemi, Finland August 29 - September 4.
- [3] P. Fayet and S. Ferrara. Supersymmetry. *Phys. Rept.*, 32:249–334, 1977.
- [4] R. Barbieri. Looking beyond the standard model: The supersymmetric option. *Riv. Nuovo Cim.*, 11N4:1–45, 1988.
- [5] S. P. Martin. *A supersymmetry primer*. Perspectives on Supersymmetry. Advanced Series on Directions in High Energy Physics - Vol 18, World Scientific, 1998.
- [6] J. A. Bagger. *Weak-scale supersymmetry: Theory and practice*. Proceedings of the Boulder 1995 TASI Lectures (QCD161:T45:1995).
- [7] H. E. Haber and G. L. Kane. The search for supersymmetry: Probing physics beyond the standard model. *Phys. Rept.*, 117:75, 1985.
- [8] G. Jungman, M. Kamionkowski, and K. Griest. Supersymmetric dark matter. *Phys. Rept.*, 267:195–373, 1996.
- [9] H. P. Nilles. Supersymmetry, supergravity and particle physics. *Phys. Rept.*, 110:1, 1984.
- [10] A. H. Chamseddine, R. Arnowitt, and P. Nath. Locally supersymmetric grand unification. *Phys. Rev. Lett.*, 49:970, 1982.

- [11] R. Barbieri, S. Ferrara, and C. A. Savoy. Gauge models with spontaneously broken local supersymmetry. *Phys. Lett.*, B119:343, 1982.
- [12] L. J. Hall, J. Lykken, and S. Weinberg. Supergravity as the messenger of supersymmetry breaking. *Phys. Rev.*, D27:2359–2378, 1983.
- [13] G. Gamberini, G. Ridolfi, and F. Zwirner. On radiative gauge symmetry breaking in the minimal supersymmetric model. *Nucl. Phys.*, B331:331–349, 1990.
- [14] J. L. Feng, K. T. Matchev, and T. Moroi. Focus points and naturalness in supersymmetry. *Phys. Rev.*, D61:075005, 2000.
- [15] J. L. Feng, K. T. Matchev, and F. Wilczek. Neutralino dark matter in focus point supersymmetry. *Phys. Lett.*, B482:388–399, 2000.
- [16] A. Djouadi, J.L. Kneur, and G. Moultaka. Suspect program, <http://www.lpm.univ-montp2.fr:7082/~kneur/suspect.html>.
- [17] A. Djouadi, M. Drees, and J. L. Kneur. Constraints on the minimal supergravity model and prospects for susy particle production at future linear e^+e^- colliders. *JHEP*, 08:055, 2001.
- [18] D. J. Castano, E. J. Piard, and Pierre Ramond. Renormalization group study of the standard model and its extensions. 2. the minimal supersymmetric standard model. *Phys. Rev.*, D49:4882–4901, 1994.
- [19] V. D. Barger, M. S. Berger, and P. Ohmann. The supersymmetric particle spectrum. *Phys. Rev.*, D49:4908–4930, 1994.
- [20] Damien M. Pierce, Jonathan A. Bagger, Konstantin T. Matchev, and Ren-jie Zhang. Precision corrections in the minimal supersymmetric standard model. *Nucl. Phys.*, B491:3–67, 1997.
- [21] P. Gondolo, J. Edsjo, L. Bergstrom, P. Ullio, and E. A. Baltz. Darksusy: A numerical package for dark matter calculations in the mssm, astro-ph/0012234, proceedings of york 2000, the identification of dark matter. pages 318–323.
- [22] P. Gondolo, J. Edsj, L. Bergstrm, P. Ullio, and T. Baltz. Darksusy program, <http://www.physto.se/~edsjo/darksusy/>.
- [23] T. Nihei, L. Roszkowski, and R. Ruiz de Austri. Towards an accurate calculation of the neutralino relic density. *JHEP*, 05:063, 2001.
- [24] B. C. Allanach. Theoretical uncertainties in sparticle mass predictions. hep-ph/0110227, submitted to aps / dpf / dpb summer study on the future of particle physics (snowmass 2001), snowmass, colorado. 30 Jun - 21 Jul 2001.
- [25] B. C. Allanach. Softsusy: A c++ program for calculating supersymmetric spectra. *Comput. Phys. Commun.*, 143:305–331, 2002.
- [26] D. E. Groom et al. Review of particle physics. *Eur. Phys. J.*, C15:1–878, 2000.
- [27] Marc Knecht and Andreas Nyffeler. Hadronic light-by-light corrections to the muon $g-2$: The pion-pole contribution. *Phys. Rev.*, D65:073034, 2002.
- [28] Andrzej Czarnecki and William J. Marciano. The muon anomalous magnetic moment: A harbinger for 'new physics'. *Phys. Rev.*, D64:013014, 2001.
- [29] J. L. Feng, K. T. Matchev, and F. Wilczek. Prospects for indirect detection of neutralino dark matter. *Phys. Rev.*, D63:045024, 2001.
- [30] S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi. Effects of supergravity induced electroweak breaking on rare b decays and mixings. *Nucl. Phys.*, B353:591–649, 1991.
- [31] M. Ciuchini, G. Degrandi, P. Gambino, and G. F. Giudice. Next-to-leading qcd corrections to $b \rightarrow \bar{c} s/\gamma$: Standard model and two-higgs doublet model. *Nucl. Phys.*, B527:21–43, 1998.
- [32] R. Barate et al. Search for the neutral higgs bosons of the standard model and the mssm in e^+e^- collisions at $s^{1/2} = 189$ -gev. *Eur. Phys. J.*, C17:223–240, 2000.
- [33] A. Heister et al. Final results of the searches for neutral higgs bosons in e^+e^- collisions at $s^{1/2}$ up to 209-gev. *Phys. Lett.*, B526:191–205, 2002.
- [34] C. Boehm, A. Djouadi, and M. Drees. Light scalar top quarks and supersymmetric dark matter. *Phys. Rev.*, D62:035012, 2000.
- [35] J. R. Ellis, K. A. Olive, and Y. Santoso. Calculations of neutralino stop coannihilation in the cmssm, cern-th-2001-339, umn-th-2032-01, tpi-minn-01-50, hep-ph/0112113.
- [36] R. Abusaidi et al. Exclusion limits on the wimp nucleon cross-section from the cryogenic dark matter search. *Phys.Rev.Lett.*, 84:5699–5703, 2000.
- [37] A. Benoit et al. Improved exclusion limits from the edelweiss wimp search, astro-ph/0206271.

- [38] G. Chardin. Edelweiss dark matter search, talk given at the school and workshop on neutrino particle astrophysics, les houches 21 jan -1st feb 2002.
- [39] N. Spooner. New limits and progress from the Boulby dark matter programme, talk given at the school and workshop on neutrino particle astrophysics, les houches 21 jan -1st feb. 2002.
- [40] F. Mayet, D. Santos, Yu. M. Bunkov, E. Collin, and H. Godfrin. Search for supersymmetric dark matter with superfluid he-3 (mache3). *Phys. Lett.*, B538:257, 2002.
- [41] Olga V. Suvorova. Status and perspectives of indirect search for dark matter, published in tegernsee 1999, beyond the desert 1999. pages 853–867.
- [42] T. Montaruli. Search for wimps using upward-going muons in macro, proceedings of the 26th icrc in salt lake city, hep-ex/9905021. pages 277–280, 17-25 Aug 1999.
- [43] A. Habis. Discriminating between $\nu/\mu \rightarrow \bar{\nu}/\bar{\mu}$, ν/τ and $\nu/\mu \rightarrow \bar{\nu}/\bar{\mu}$ (sterile) in atmospheric ν/μ oscillations with the super-kamiokande detector, proceedings of the 27th icrc, hamburg, germany, hep-ex/0106024. 7-15 Aug 2001.
- [44] L. Thompson. Dark matter prospects with the antares neutrino telescope, talk given at the conference dark 2002, cape town, south africa 4-9 feb. 2002.
- [45] J. Edsjo. Swedish astroparticle physics, talk given at the conference 'partikeldagarna', uppsala, sweden, march 6. 2001.
- [46] J. R. Ellis, T. Falk, K. A. Olive, and M. Srednicki. Calculations of neutralino stau coannihilation channels and the cosmologically relevant region of mssm parameter space. *Astropart. Phys.*, 13:181–213, 2000.
- [47] V. Bertin, E. Nezri, and J. Orloff. in preparation.
- [48] Lars Bergstrom, Joakim Edsjo, and Paolo Gondolo. Indirect detection of dark matter in km-size neutrino telescopes. *Phys. Rev.*, D58:103519, 1998.
- [49] John R. Ellis, Andrew Ferstl, and Keith A. Olive. Re-evaluation of the elastic scattering of supersymmetric dark matter. *Phys. Lett.*, B481:304–314, 2000.
- [50] John R. Ellis, Andrew Ferstl, and Keith A. Olive. Constraints from accelerator experiments on the elastic scattering of cmssm dark matter. *Phys. Lett.*, B532:318–328, 2002.
- [51] Vernon D. Barger, Francis Halzen, Dan Hooper, and Chung Kao. Indirect search for neutralino dark matter with high energy neutrinos. *Phys. Rev.*, D65:075022, 2002.
- [52] L. Roszkowski, R. Ruiz de Austri, and T. Nihei. New cosmological and experimental constraints on the cmssm. *JHEP*, 08:024, 2001.
- [53] Howard Baer, Frank E. Paige, Serban D. Protopopescu, and Xerxes Tata. Isajet 7.48: A monte carlo event generator for p p, anti-p p, and e+ e- reactions, bnl-het-99-43, fsu-hep-991218, uh-511-952-00, hep-ph/0001086. 1999.